



Changing medication method influences total amount of administered antimicrobials

Fertner, Mette Ely; Boklund, Anette ; Toft, Nils

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Fertner, M. E., Boklund, A., & Toft, N. (2015). *Changing medication method influences total amount of administered antimicrobials*. Poster session presented at SVEPM Annual Meeting 2015, Ghent, Belgium.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Changing medication method influences total amount of administered antimicrobials

A register based study

Mette Fertner, Anette Boklund, Nils Toft
Section for Epidemiology, National Veterinary Institute, Technical University of Denmark




Total amount of antimicrobials prescribed for Danish finishers during 2005-2013, by administration form.

Conclusion


Finisher herds going from feed- to water-administered antimicrobials for group treatment during 2008-09 significantly increased their total use of antimicrobials by 1.33 [0.33; 2.33] ADD₅₀/100 finishers/day. A general change of medication method may explain some of the increased use of antimicrobials during 2008-09.

Data & Results



For all three groups, the difference in prescribed antimicrobials between 2008-09 was calculated. Subsequently, it was tested whether the difference between years varied significantly between the three study groups.

Test	p-value
Kruskal-Wallis	0.0001
Pairwise comparison	
feed/water – feed/feed	0.017
feed/water – water/water	0.037
feed/feed – water/water	0.709



Use of antimicrobials for finishers during 2008-09 in the three study groups.

